



## **Transient increase in neuronal chloride concentration by neuroactive aminoacids released from glioma cells.**

Cristina Bertollini, Emanuele Murana, Luciana Mosca, Maria d'Erme, Federico Scala, Antonio Francioso, Myriam Catalano, Cristina Limatola, Piotr Bregestovski, Silvia Di Angelantonio, et al.

### **► To cite this version:**

Cristina Bertollini, Emanuele Murana, Luciana Mosca, Maria d'Erme, Federico Scala, et al.. Transient increase in neuronal chloride concentration by neuroactive aminoacids released from glioma cells.. Frontiers in Molecular Neuroscience, 2012, 5, pp.100. 10.3389/fnmol.2012.00100 . pasteur-00961590

**HAL Id: pasteur-00961590**

**<https://hal-riip.archives-ouvertes.fr/pasteur-00961590>**

Submitted on 20 Mar 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Transient increase in neuronal chloride concentration by neuroactive aminoacids released from glioma cells

Cristina Bertollini<sup>1\*</sup>, Emanuele Murana<sup>1</sup>, Luciana Mosca<sup>2</sup>, Maria D'Erme<sup>2</sup>, Federico Scala<sup>1</sup>, Antonio Francioso<sup>2</sup>, Myriam Catalano<sup>1,3</sup>, Cristina Limatola<sup>1,3</sup>, Piotr Bregestovski<sup>4</sup>, Silvia Di Angelantonio<sup>1,5†</sup> and Davide Ragozzino<sup>1,3†</sup>

<sup>1</sup> Department of Physiology and Pharmacology, Istituto Pasteur-Fondazione Cenci Bolognetti, Sapienza University of Rome, Rome, Italy

<sup>2</sup> Department of Biochemical Sciences, Sapienza University of Rome, Rome, Italy

<sup>3</sup> IRCCS Neuromed, Pozzilli, Italy

<sup>4</sup> INSERM URM 1106, Aix-Marseille Université Brain Dynamics Institute, Marseille, France

<sup>5</sup> Center for Life Nano Science at Sapienza, Istituto Italiano di Tecnologia—Sapienza University of Rome, Rome, Italy

## Edited by:

Pau Gorostiza, Institutió Catalana de Recerca i Estudis Avançats, Spain

## Reviewed by:

Rafael Gutierrez, Centro de Investigación y Estudios Avanzados del IPN, Mexico  
Daniele Arosio, Consiglio Nazionale delle Ricerche, Italy

## \*Correspondence:

Cristina Bertollini, Department of Physiology and Pharmacology, Fondazione Cenci Bolognetti, Sapienza University of Rome, Piazzale Aldo Moro 5, Rome, Italy.  
e-mail: cristina.bertollini@uniroma1.it

<sup>†</sup> These authors equally contributed to this work.

Neuronal chloride concentration ( $[Cl^-]_i$ ) is known to be dynamically modulated and alterations in  $Cl^-$  homeostasis may occur in the brain at physiological and pathological conditions, being also likely involved in glioma-related seizures. However, the mechanism leading to changes in neuronal  $[Cl^-]_i$  during glioma invasion are still unclear. To characterize the potential effect of glioma released soluble factors on neuronal  $[Cl^-]_i$ , we used genetically encoded CFP/YFP-based ratiometric  $Cl^-$  (apical) Sensor transiently expressed in cultured hippocampal neurons. Exposition of neurons to glioma conditioned medium (GCM) caused rapid and transient elevation of  $[Cl^-]_i$ , resulting in the increase of fluorescence ratio, which was strongly reduced by blockers of ionotropic glutamate receptors APV and NBQX. Furthermore, in HEK cells expressing GluR1-AMPA receptors, GCM activated ionic currents with efficacy similar to those caused by glutamate, supporting the notion that GCM contains glutamate or glutamatergic agonists, which cause neuronal depolarization, activation of NMDA and AMPA/KA receptors leading to elevation of  $[Cl^-]_i$ . Chromatographic analysis of the GCM showed that it contained several aminoacids, including glutamate, whose release from glioma cells did not occur via the most common glial mechanisms of transport, or in response to hypoosmotic stress. GCM also contained glycine, whose action contrasted the glutamate effect. Indeed, strychnine application significantly increased GCM-induced depolarization and  $[Cl^-]_i$  rise. GCM-evoked  $[Cl^-]_i$  elevation was not inhibited by antagonists of  $Cl^-$  transporters and significantly reduced in the presence of anion channels blocker NPPB, suggesting that  $Cl^-$  selective channels are a major route for GCM-induced  $Cl^-$  influx. Altogether, these data show that glioma released aminoacids may dynamically alter  $Cl^-$  equilibrium in surrounding neurons, deeply interfering with their inhibitory balance, likely leading to physiological and pathological consequences.

**Keywords:**  $Cl^-$ -Sensor, glioma cells, hippocampus, glutamate, anionic channels

## INTRODUCTION

In the central nervous system (CNS), a tight regulation of intracellular chloride concentration ( $[Cl^-]_i$ ) is important for a number of cellular functions, including the stabilization of resting membrane potential, the regulation of both intracellular pH and

cell-volume (Pasantes-Morales et al., 2006; Suzuki et al., 2006) and the strength and polarity of  $\gamma$ -aminobutyric acid (GABA) and glycine-mediated neurotransmission (Payne et al., 2003).

Several different pathways allow  $Cl^-$  movements across neuronal membranes, determining  $Cl^-$  equilibrium. Among these are the ligand-gated anion channels (GABA<sub>A</sub> and glycine receptors), the cation-chloride cotransporters (KCC2 and NKCC1) and a variety of  $Cl^-$  channels, including  $Ca^{2+}$ -, volume-, and voltage-activated  $Cl^-$  channels (Payne et al., 2003; Suzuki et al., 2006; Jentsch, 2008; Deisz et al., 2011).

Neuronal  $Cl^-$  equilibrium is subject to alterations both in physiological and pathological conditions (Staley et al., 1995; Planells-Cases and Jentsch, 2009; Doyon et al., 2011) and abnormal  $Cl^-$  homeostasis is associated with neuronal trauma or brain disorders (De Koninck, 2007). Particularly, in epilepsy, the alteration of  $Cl^-$  homeostasis is a widespread phenomenon,

**Abbreviations:** ACM, Astrocyte conditioned medium; APV, D-2-amino-5-phosphonopentanoic;  $[Cl^-]_i$ , chloride concentration; CNS, central nervous system; DMEM, Dulbecco's Modified Eagle Medium; DMSO, Dimethyl sulfoxide; EAATs, Excitatory aminoacid transporters; FBS, Fetal Bovine Serum; FFA, flufenamic acid; GABA,  $\gamma$ -aminobutyric acid; GCM, glioma conditioned medium; Glu, glutamate; GluR1-HEK cells, HEK 293 stably expressing the rat flip variant of wild-type glutamate receptor 1; HBSS, Hank's balanced salt solution; HPLC, High Performance Liquid Chromatography; HyperGCM, hyperosmotic glioma conditioned medium; NBQX, 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione; NES, normal external solution; NFA, niflumic acid; NPPB, 5-Nitro-2-(3-phenylpropylamino)benzoic acid; OPA, *o*-phthalaldehyde; SAS, Sulfasalazine; TBOA, DL-threo- $\beta$ -Benzoyloxyaspartic acid; TTX, Tetrodotoxin.

contributing to neuronal hyperexcitability (Palma et al., 2006; Barmashenko et al., 2011; Conti et al., 2011). The reported changes in neuronal  $\text{Cl}^-$  equilibrium affect the inhibitory power of GABA/glycine, causing its switching to excitatory action. In addition, neuronal transmembrane  $\text{Cl}^-$  gradient may be altered by glutamatergic stimulation, through the activation of anionic channels or changes in  $\text{Cl}^-$  transporters activity (Van Damme et al., 2003; Slemmer et al., 2004; Kitamura et al., 2008). In the brain, a similar condition may occur in case of excessive glutamate release, under intense synaptic activity (Fiumelli et al., 2005; Fiumelli and Woodin, 2007) or reduced glutamate uptake by astrocytes (Danbolt, 2001). Indeed, even modest increases in extracellular glutamate concentration can alter synaptic transmission (Araque et al., 1999) and reduced activity of glial glutamate transporters has been suggested to contribute to and exacerbate a number of neurological conditions, including stroke, epilepsy, cerebral ischaemia, amyotrophic lateral sclerosis, and others (O'Shea, 2002).

A dysfunction in glutamate transport has also been reported in malignant gliomas (Ye et al., 1999). The comparison of glutamate transport into astrocytes vs. their malignant counterparts showed that gliomas behave opposite to astrocytes, releasing glutamate rather than sequestering it (Ye and Sontheimer, 1999). Moreover, when glioma cells were co-cultured with neurons, the released glutamate activated neuronal NMDA receptors, resulting in excitotoxic cell death, suggesting that the release of excitotoxic concentrations of glutamate may promote tumor expansion (Sontheimer, 2003). In addition, glutamate, as well as other aminoacids (GABA, glycine, serine) may be released, in the brain, by various cell types during volume changes (Pasantes-Morales et al., 2006), occurring upon cell migration, edema, or during glioma cell invasion (Ordaz et al., 2004; Pasantes-Morales and Vázquez-Juárez, 2012).

In this study we analyzed how glutamate or other aminoacids, released by glioma cells cause an alteration of  $\text{Cl}^-$  homeostasis. Taking advantage of a genetically encoded CFP/YFP-based ratio-metric chloride sensor (Cl-Sensor) (Markova et al., 2008; Waseem et al., 2010), we investigated how neuronal  $[\text{Cl}^-]_i$  is dynamically regulated by diffusible factors released by cultured glioma cells.

We report that glutamate released from glioma cells triggers neuronal  $\text{Cl}^-$  rise through the activation of ionotropic glutamate receptors, cell depolarization and activation of anionic channels. Such mechanism may partially explain how glioma invasion causes local dynamic  $\text{Cl}^-$  changes and neuronal  $\text{Cl}^-$  equilibrium dysfunction.

## MATERIALS AND METHODS

### ANIMALS

Animal procedures were conducted in accordance with the international guidelines on the ethical use of animals from the European Communities Council Directive of 24 November 1986 (86/609/EEC). C57BL/6 (Charles River Laboratory) of either sex was used.

### CELL CULTURES

The glioblastoma patient derived cell line MZC (M.Z.C., kindly provided by Dr. Antonietta Arcella, Neuromed, Italy, Sciacaluga

et al., 2010) was grown in Dulbecco's Modified Eagle Medium (DMEM, GIBCO) supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 100 IU/ml penicillin G and 100  $\mu\text{g}/\text{ml}$  streptomycin at 37°C in a 5%  $\text{CO}_2$  humidified atmosphere. Medium was changed twice a week and the cells were sub-cultivated when confluent and used between 20th and 40th passage when they had reached 80% confluence.

Primary hippocampal neuronal cultures were obtained from postnatal day 0–2 (P0–P2) C57BL/6 mice. Briefly, after careful dissection from diencephalic structures, the meninges were removed under a dissection microscope and the hippocampi were collected in ice-cold Hank's balanced salt solution (HBSS, GIBCO Invitrogen) and chopped and digested in 1.25 mg/ml trypsin for 15 min at 37°C. Subsequently, cells were mechanically dissociated and plated at a density of  $10^5$  in poly-L-lysine coated round glass coverslips (12 mm diameter) put into 24-well culture plates in serum-free Neurobasal medium, supplemented with 2% B27, 0.5 mM L-Glutamine and 100  $\mu\text{g}/\text{ml}$  gentamicin (culture medium). Finally, cells were kept at 37°C in 5%  $\text{CO}_2$  for 10–12 days with a twice a week medium replacement (1:1 ratio). With this method we obtained 60–70% neurons, 30–35% astrocytes, and 4–5% microglia, as determined with  $\beta$ -tubulin III, glial fibrillary acidic protein, and isolectin IB4 staining (Lauro et al., 2010).

Primary cortical glial cells were prepared from P0–P2 -old mice. Cerebral cortices were chopped and digested in 30 U/ml papain for 40 min at 37°C followed by gentle trituration. The dissociated cells were washed, suspended in DMEM with 10% FBS (Invitrogen) and 2 mM l-glutamine and plated at a density of  $9\text{--}10 \times 10^5$  in 175  $\text{cm}^2$  cell culture flasks. At confluence (10–14 DIV), glial cells were shaken for 2 h at 37°C to remove microglial cells. These procedures gave almost pure astrocytes cell population (4–6% of microglia contamination), as verified by staining with glial fibrillary acidic protein and isolectin IB4 (Rosito et al., 2012). Astrocytes were re-plated and used for experiments.

The human retroviral packaging cell line HEK 293 stably expressing the rat flip variant of wild-type glutamate receptor 1 (GluR1-HEK cells) was grown in DMEM with geneticin (0.5 mg/ml), Glutamax-I/10% FBS 1% penicillin/streptomycin, 5%  $\text{CO}_2$  (37°C). Cells were plated onto 12 mm round coverslips at a density of 104 cells/ $\text{cm}^2$ .

### GLIOMA CONDITIONED MEDIUM (GCM)

Confluent MZC cultures were incubated for 4 h with 10 ml of filtered normal external solution (NES) containing (in mM): 140 NaCl, 2.8 KCl, 2  $\text{CaCl}_2$ , 2  $\text{MgCl}_2$ , 10 HEPES-NaOH, and 10 glucose (pH 7.32;  $300 \pm 5$  mOsm). This GCM was then centrifuged, pH adjusted to 7.32 and used for the experiments. The hyperosmotic glioma conditioned medium (HyperGCM) was obtained by incubating glioma cells with the following external solutions in (mM): NaCl 159.6, KCl 3.2,  $\text{MgCl}_2$  2.28,  $\text{CaCl}_2$  2.28, HEPES-NaOH 11.4, glucose 11.4 (114% concentrated NES); or NaCl 154, KCl 2.8,  $\text{MgCl}_2$  2,  $\text{CaCl}_2$  2, HEPES-NaOH 10, glucose 25. Both solutions, adjusted at pH 7.32, were isoosmotic compared to DMEM ( $340 \pm 10$  mOsm).

## HIPPOCAMPAL CULTURES TRANSFECTION

Twenty Four hours before transfection, 50% of the neuronal growth medium was replaced with fresh culture medium. For transfection 100  $\mu\text{l}$  of MEM was mixed with 2  $\mu\text{l}$  of Magnetofection NeuroMag (OZ Bioscience, France) and 1  $\mu\text{g}$  of the cDNA of Cl-Sensor (Markova et al., 2008). The mixture was incubated for 15–20 min at room temperature and thereafter distributed dropwise over the neuronal culture. Cells were then placed on a magnetic plate (OZ Bioscience) and incubated for 15 min at 37°C. Transfection was terminated by substitution of 50% of incubation solution with the fresh culture medium. Cells were used for the experiments 24–76 h after transfection.

## FLUORESCENCE DETERMINATIONS

Fluorescence images were acquired at room temperature (22–24°C) using a customized digital imaging microscope. Excitation of cells at various wavelengths was achieved using a 1-nm-bandwidth polychromatic light selector (Till Polychrome V) equipped with a 150 W xenon lamp (Till Photonics, Germany). For the use of Cl-Sensor the optic system is equipped with an excitation filter D480/30x, an emission filter D535/40m and a dichroic mirror 500DRLP (500 nm) (Omega Optics, USA); for the use of FURA-2AM we used a conventional set of filters (SP410, 510/40 m; 400DCLP 400 nm; Omega Optics, USA). Fluorescence was visualized using an upright microscope (Axioskope) equipped with a 40  $\times$  water-immersion objective (Achromplan CarlZeiss, USA) and a digital 12 bit CCD camera system (SensiCam, PCO AG, Germany); images were acquired on a computer via a double-coaxial cable. All peripheral hardware control, image acquisition and image processing were achieved using customized software TillVision v. 4.0 (Till Photonics, Germany).

Cells expressing Cl-Sensor were excited alternatively at 445 and 485 nm wavelengths (50 ms, 0.1 Hz).  $[\text{Cl}^-]_i$  changes are expressed as a ratio of background subtracted  $F_{445}$  over  $F_{485}$  ( $R = F_{445}/F_{485}$ ). Before starting all the experiments, we evaluated the basal  $\text{Cl}^-$  level in each neuron, through the emission spectrum (for methods see Markova et al., 2008; Bregestovski et al., 2009). Cells whose shape of emission spectrum corresponded to high  $[\text{Cl}^-]_i$  were not considered.

The  $\text{Ca}^{2+}$  sensitive indicator Fura-2 was used to monitor changes in the intracellular calcium level in hippocampal neurons. Cells (10–12 DIV) were loaded by adding cell permeant acetoxymethyl (AM) derivatives dye (4  $\mu\text{M}$ ) and incubating dishes at 37°C for 45 min. Fura-2AM loaded cells were excited alternatively with dual wavelengths of 340 and 380 nm (20 ms; 0.2 Hz). Intracellular  $\text{Ca}^{2+}$  changes are expressed as a ratio between fluorescence emitted after excitation at 340 nm ( $F_{340}$ ) and after excitation at 380 nm ( $F_{380}$ ) [Fluorescence Ratio ( $R$ ) =  $F_{340}/F_{380}$ ].

## PATCH-CLAMP RECORDINGS

Whole-cell patch-clamp recordings were performed using a Axopatch 200A amplifier, Digidata 1440 digitizer, on a PC running pCLAMP software (all from Molecular Devices, Foster city, CA). Data were acquired at 10 kHz and filtered at 2 kHz. Patch electrodes were prepared from borosilicate glass using a

vertical electrode puller (PC10, Narishige, Japan) to produce tip openings of 1–2  $\mu\text{m}$  (4–6 M $\Omega$ ). Electrodes were filled with an intracellular solution containing (in mM): 135 CsMetSO<sub>4</sub>, 2 MgCl<sub>2</sub>, 2 MgATP, 0.3 NaGTP, 10 HEPES, 0.5 EGTA; pH 7.3 with CsOH. Recording were performed in normal extracellular solution containing (in mM): 140 NaCl, 2.5 KCl, 2 CaCl<sub>2</sub>, 2 MgCl<sub>2</sub>, 10 HEPES–NaOH, and 10 glucose (pH 7.3 adjusted with NaOH, osmolarity 300 mOsm). In voltage-clamp mode of recording cells were held at –70 mV; in current-clamp experiments, the voltage was recorded while cells were injected with holding current (30–230) pA. Glutamate concentration–current curves were constructed by applying to each HEK-AMPA cell (held at –50 mV) four or five different concentrations of agonist (0.01–1 mM) at 60–120 s intervals, in the presence of 25  $\mu\text{M}$  cyclothiazide, to prevent GluR1 receptor desensitization (Fucile et al., 2006), and normalizing the current response to the plateau value, tested in all cells (corresponding to 1 mM). For each cell, the concentration–current curve was best-fitted using Origin 7 (OriginLab Corp., Northampton, MA, USA) to the Hill equation:  $I_{\text{norm}} = 1/(1 + \text{EC}_{50}^{n_H}/[\text{Glutamate}]^{n_H})$  where  $I_{\text{norm}}$  is the normalized current response,  $\text{EC}_{50}$  is the agonist concentration yielding half-maximal current response and  $n_H$  is the Hill coefficient. These curves were averaged and used to calculate the mean  $\text{EC}_{50}$  for glutamate.

## EXPERIMENTAL PROCEDURES AND DRUGS APPLICATION

During fluorescence or electrophysiological measurements, cells were continuously superfused with external solution, using a gravity driven perfusion system, positioned 50–100  $\mu\text{m}$  from the cell. Unless otherwise indicated, all experiments were conducted in normal extracellular solution. Concentration response curves in HEK-AMPA cells were performed with a computer controlled fast perfusion system (Warner Instruments, USA). Tetrodotoxin (TTX, 1  $\mu\text{M}$ ) was routinely dissolved in the extracellular medium. Low  $\text{Cl}^-$  medium contained: 147.5 mM Na-Gluconate, 2.8 mM K-Gluconate, 2 mM MgCl<sub>2</sub>, 2 mM CaCl<sub>2</sub>, 10 mM Glucose, and 10 mM HEPES–NaOH (pH 7.32; 300  $\pm$  5 mOsm). The fluorescence signal was monitored online and stable baseline responses were recorded for at least 4 min before applying drugs. Tetrodotoxin citrate (TTX), D-2-amino-5-phosphonopentanoic (D-APV), 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione (NBQX), glutamate, GABA, strychnine hydrochloride, and bicuculline methochloride were purchased from Abcam (UK). 5-Nitro-2-(3-phenylpropylamino)benzoic acid (NPPB) and DL-threo- $\beta$ -Benzyloxyaspartic acid (TBOA) were purchased from Tocris (UK). All other drugs used were purchased from Sigma-Aldrich, Milan, Italy.

NPPB, Sulfasalazine (SAS), and TBOA were dissolved in Dimethyl sulfoxide (DMSO) and used at 1/1000 dilution and the same concentration of solvent was used in control condition. Unless otherwise indicated, AP5, NBQX, and strychnine were applied to the cells only in co-application with GCM. The duration of NPPB pretreatment (3–4 min) was set to allow stabilization of fluorescence ratio after the change induced by drug application.



## CHROMATOGRAPHIC ANALYSIS

Aminoacids were determined in supernatants by reversed-phase High Performance Liquid Chromatography (HPLC) following *o*-phthaldialdehyde (OPA) derivatization in the presence of 2-mercaptoethanol. 100  $\mu\text{L}$  of samples or standards were mixed 1:1 (v/v) with freshly prepared OPA/2-mercaptoethanol derivatizing solution (3,5 mg of OPA in 50  $\mu\text{L}$  of 95% ethanol, diluted with 5 mL of borate buffer, pH 10.4, plus 10  $\mu\text{L}$  of 2-mercaptoethanol, protected from light) and injected within 1 min onto the column (Hirschberger et al., 1985). The HPLC consisted of a Waters apparatus equipped with a 600 pump and pump controller, a Rheodyne injection valve with a 50  $\mu\text{L}$  loop, a Nova-Pack C18 column (reverse phase,  $3.9 \times 150$  mm, 4  $\mu\text{m}$  particle size, thermostated at  $41^\circ\text{C}$  with a 10 mm guard column of the same material matrix) and a Shimadzu RF-551 fluorometric detector, operating at  $\lambda_{\text{ex}}$  360 nm and  $\lambda_{\text{em}}$  455 nm. The elution was performed at a flow rate of 1 mL/min, with solvent A being 25 mM phosphate buffer, pH 7.0, containing 3% THF, and solvent B 25 mM phosphate buffer, pH 7.0, containing 40%  $\text{CH}_3\text{CN}$  and 3% THF. The mobile phase was run isocratically at 5% B for the first 5 min, then solvent B linearly increased to 27% in 10 min, remained for 20 min at 27% B and finally increased linearly to 100% in 1 min. At 40 min, buffer B was returned to 5% and the column was allowed to equilibrate for 10 min. Total run time was 50 min per sample. Peak identification was performed on the basis of the retention time and by spiking the sample with appropriate standards, whereas aminoacid quantitation was performed by automatic peak area integration using dedicated software (Millennium<sup>32</sup>, Waters). Results are expressed as  $\mu\text{moles}$  of aminoacid per L of sample.

## DATA ANALYSIS AND STATISTICS


Data are analyzed with Clampfit 10 (Molecular Devices). Statistical significance ( $p < 0.05$ ) was determined using the paired or unpaired Student's *t*-test (Origin software; Microcal Software, Northampton, MA). Data are reported as means  $\pm$  standard error of the mean (SEM) and "*n*" values refers to the number of neurons examined. All studies involved different culture sets. Each set of experiments presented was internally controlled, using similar numbers of control coverslips to compare with each experimental condition.

## RESULTS

### GCM EVOKES RAPID AND REVERSIBLE INCREASE IN $[\text{Cl}^-]_i$ VIA A GLUTAMATERGIC MECHANISM IN CULTURED HIPPOCAMPAL NEURONS


In order to assess whether substances released from glioma cells may alter neuronal chloride homeostasis, we used a CFP/YFP-based ratiometric Cl-Sensor expressed in primary hippocampal neuronal cultures (Figure 1A). A glioma conditioned medium (GCM), prepared exposing cultured MZC glioma cells to NES, was applied to neurons transfected with Cl-Sensor and changes in fluorescence ratio (*R*), associated to  $[\text{Cl}^-]_i$  variations, were monitored over time.

Results indicate that GCM applied for 1 min onto neurons expressing Cl-Sensor evokes a rapid and reproducible increase in fluorescence ratio (*R*) from  $0.63 \pm 0.02$  to  $0.94 \pm 0.04$



( $\Delta R = 0.32 \pm 0.03$ ,  $n = 153$ ,  $p < 0.01$ ), indicating a strong rise in  $[\text{Cl}^-]_i$  (Figure 1B). *R* increase was followed by a slow recovery of basal fluorescence, with an average half decay time (*t*/2) of  $3.5 \pm 0.3$  min ( $n = 123$ ). As shown in Figure 1C, the duration of *t*/2 increased with peak amplitude of response to GCM.

Since glioma cells may release glutamate in the extracellular space (Ye and Sontheimer, 1999; Buckingham et al., 2011) we wondered whether GCM-induced neuronal  $[\text{Cl}^-]_i$  rise could be triggered by the activation of ionotropic glutamate receptors on neurons (see Slemmer et al., 2004). To address this issue, we treated hippocampal neurons with NMDA and AMPA receptors antagonists (APV 20  $\mu\text{M}$ ; NBQX 10  $\mu\text{M}$ ; 3 min), before exposing them to GCM. As shown in Figure 2A, APV and NBQX strongly reduced GCM-induced responses, giving an average decrease of  $\Delta R$  by  $81 \pm 8\%$  of control ( $n = 5$ ,  $p < 0.01$ ), indicating that GCM leads to a rapid increase of  $[\text{Cl}^-]_i$ , by a glutamatergic mechanism. Consistently, the exposure of Cl-Sensor expressing neurons to glutamate (100  $\mu\text{M}$ ) for 1 min evoked a rapid and reversible increase in *R* from  $0.56 \pm 0.02$  to  $1.01 \pm 0.09$  ( $\Delta R = 0.45 \pm 0.08$ ,  $n = 28$ ,  $p < 0.01$ ), similar to that induced by the application of GCM (Figure 2B). It has been reported that glutamate can cause changes both in  $[\text{Cl}^-]_i$  and  $[\text{pH}]_i$  in neurons (Wang et al., 1994; Metzger et al., 2002) and Cl-Sensor is known to be sensitive to pH (Markova et al., 2008). Thus, we monitored



**FIGURE 2 | Glutamatergic mechanism GCM-evoked  $[\text{Cl}^-]_i$  increase. (A)** Effect of ionotropic glutamate receptors antagonists on GCM-evoked  $[\text{Cl}^-]_i$  increase. Left, fluorescence trace from an example neuron showing the effect of application of GCM alone and in the presence of APV (20  $\mu\text{M}$ ) and NBQX (10  $\mu\text{M}$ ). Right, bar chart represents the average of fluorescence response elicited by GCM in the presence of APV (20  $\mu\text{M}$ ) and NBQX (10  $\mu\text{M}$ ) expressed as the % of control ( $n = 5$  neurons). **(B)** Time course of the effect of 100  $\mu\text{M}$  glutamate on the fluorescence ratio in Cl-Sensor transfected hippocampal neurons ( $n = 28$ ). **(C)** Fluorescence response to glutamate in control and in gluconate-based low  $\text{Cl}^-$  medium (8 mM); Left, fluorescence trace from an example neuron showing the effect of application of glutamate

(100  $\mu\text{M}$ ) in control and in gluconate-based low  $\text{Cl}^-$  medium (8 mM). Right, bar chart represents the average of fluorescence response elicited by glutamate in the absence of external  $\text{Cl}^-$  expressed as the % of control ( $n = 6$  neurons). **(D)** Concentration-current response curve for glutamate obtained in HEK-AMPA cells (filled squares,  $\blacksquare$ ;  $n = 12$ ). Empty circle ( $\circ$ ) represents the average amplitude of the current evoked by GCM application in the same cells, indicative of a glutamate concentration in GCM of  $20 \pm 2 \mu\text{M}$ . ( $n = 6$ ). Inset, typical whole-cell currents evoked by glutamate (10–30–100  $\mu\text{M}$ , left) and GCM (right) on HEK-AMPA cells **(A–D)**, horizontal bars represents drugs application as indicated. Data are shown as mean values  $\pm$  SEM.  $**p < 0.01$ .

fluorescence response to glutamate in control and in gluconate-based low  $\text{Cl}^-$  medium (8 mM, similar to the one measured in Cl-Sensor expressing neurons), to exclude that the fluorescence transients could depend on the effect of intracellular acidification on Cl-Sensor properties. The removal of external  $\text{Cl}^-$  abolished  $R$  increase in response to glutamate ( $\Delta R = 0.03 \pm 0.01$ ,  $n = 6$ ,  $p < 0.05$ ), suggesting that, in our experimental conditions, the glutamate-induced fluorescence ratio increase truly represents  $[\text{Cl}^-]_i$  rise (Figure 2C).


Then, in order to functionally evaluate the concentration of glutamate present in GCM, we constructed a concentration-current response curve for glutamate in HEK cell lines stably expressing the GluR1 subunit of AMPA receptors (HEK-AMPA cells) (Fucile et al., 2000), using five different glutamate concentrations and applying GCM on the same cell. Best fits of concentration-current response curves were averaged and the

mean  $\text{EC}_{50}$  for glutamate was  $59.4 \pm 0.6 \mu\text{M}$  ( $n_H = 1.30 \pm 0.01$ ,  $n = 24$ ). The amplitude of currents evoked by GCM application in the same cells were fitted in the curve, indicating a glutamate concentration of  $20 \pm 2 \mu\text{M}$  (Figure 2D  $n = 6$ ). Consistently, the application of 20  $\mu\text{M}$  glutamate to neurons transfected with Cl-Sensor was sufficient to cause a transient increase in  $[\text{Cl}^-]_i$  ( $\Delta R = 0.06 \pm 0.01$ ,  $n = 3$ ; not shown).

Altogether, these data suggest that glutamate released from glioma cells triggers neuronal  $\text{Cl}^-$  rise through the activation of ionotropic glutamate receptors.

#### GLIOMA CONDITIONED MEDIUM CONTAINS EXCITATORY AND INHIBITORY AMINOACIDS

We used the High-Performance Liquid Chromatography (HPLC) to quantify the amount of glutamate contained in GCM and identify other aminoacids eventually present. Figure 3A shows



a representative chromatogram of the aminoacids contained in GCM. The HPLC analysis confirmed that glioma cells release a significant amount of glutamate, in addition to several other aminoacids in the micromolar range, including alanine, aspartate, serine, and glycine (Behrens et al., 2000). **Table 1** summarizes the concentration of the aminoacids present in the GCM. Among these aminoacids, we highlighted both excitatory (glutamate, aspartate, glycine, serine) and inhibitory (glycine, taurine) neuroactive compounds. In particular, glutamate seems to be the most relevant excitatory neurotransmitter, with an average concentration in GCM of  $10.6 \pm 1.7 \mu\text{M}$  ( $n = 11$ ). On the other hand, among inhibitory neurotransmitters, glycine is by far the most representative, as its concentration in GCM is appreciably above the level of taurine or GABA, the latter being undetectable.

Since under specific/pathological circumstances, astrocytes may also release glutamate, (Malarkey and Parpura, 2008), we performed HPLC analysis on astrocytes-conditioned medium, observing an extremely low concentration of glutamate

**Table 1 | Concentration of identified aminoacids in glioma and astrocyte-conditioned medium; table summarizes the concentration of aminoacids determined in GCM ( $n = 11$ ) and astrocyte conditioned medium (ACM) ( $n = 4$ ) expressed in  $\mu\text{M}$ .**

AA	Concentration in GCM ( $\mu\text{M}$ )	Concentration in ACM ( $\mu\text{M}$ )
Aspartate	$2.3 \pm 0.3$	$6.7 \pm 4.1$
Glutamate	$10.6 \pm 1.7$	$1.5 \pm 0.9$
Serine	$8.5 \pm 1.4$	$31.6 \pm 10.1$
Glycine	$16.0 \pm 2.7$	$22.3 \pm 6.0$
Alanine	$51.7 \pm 7.2$	$32.0 \pm 4.0$
Taurine	$1.0 \pm 0.5$	$4.5 \pm 1.4$

Glutamate was not quantifiable in 3/3 astrocyte conditioned medium tested; Alanine was detected only in 1/3 astrocyte conditioned medium at  $32 \mu\text{M}$ .

( $1.5 \pm 0.9 \mu\text{M}$ ,  $n = 4$ ) in respect to GCM. These data indicate that, in our experimental conditions, glutamate release is a specific feature of glioma cells.

Trying to identify the mechanism of glutamate release from glioma cells, we first put attention to the  $X_c$  transport system (Buckingham et al., 2011), analyzing the GCM obtained in the presence of the  $X_c$  system blocker sulfasalazine (SAS,  $250 \mu\text{M}$ , 4 h). As shown in **Figure 3B**, SAS treatment did not reduce the amount of glutamate released in the GCM ( $94 \pm 10\%$  of control;  $n = 4$ ). We further tried to interfere with glutamate release from glioma cells, by blocking glutamate transporters with the competitive, non-transportable excitatory aminoacid transporters (EAATs) blocker  $\beta$ -threo-benzoyloxy aspartate (TBOA, Shimamoto et al., 1998). However, GCM prepared in the presence of TBOA ( $100 \mu\text{M}$ ), showed an increased glutamate level ( $148 \pm 16\%$ ,  $p = 0.01$ ;  $n = 4$ ), indicating that in MZC cells EAAT transport is not necessary for glutamate release and rising the suggestion that it may be involved in glutamate uptake (Takano et al., 2001).

Another possible way of glutamate release is linked to osmotic stress (Takano et al., 2005). Since the extracellular solution used for the preparation of GCM, is hypotonic (NES,  $300 \pm 5 \text{ mOsm}$ ) compared to glioma culture medium ( $340 \pm 10 \text{ mOsm}$ ; DMEM), glutamate could be released in response to an osmotic stress. Indeed, it has been reported that several aminoacids are released from neuronal and non-neuronal cells in response to hypotonic stress (Ordaz et al., 2004; Takano et al., 2005; Pasantes-Morales and Vázquez-Juárez, 2012). In particular, glial cells are able to respond to hypoosmotic swelling, with a regulatory volume decrease, through the efflux of chloride, glutamate, and other anions (Kimelberg et al., 1995; Pasantes-Morales and Vázquez-Juárez, 2012). To test this possibility, we analyzed the level of glutamate in the GCM, prepared with a hyperosmotic solution (HyperGCM), having the same osmolality of the glioma culture medium ( $\sim 340 \text{ mOsm}$ ). The concentration of glutamate determined in the HyperGCM by HPLC analysis was  $12.5 \pm 2.7 \mu\text{M}$  (**Figure 3C**;  $n = 5$ ), similar to that measured in the GCM, suggesting that the osmotic stress is not involved in glutamate release. The same holds for all the identified aminoacids (with the exception of taurine),

whose concentrations were similar in the two conditions (**Figure 3C**).

Altogether, these data indicate that glioma cells release both excitatory and inhibitory neuroactive compounds. Among them, glutamate is released, independently of the most conventional release mechanisms, from glioma cells but not from astrocytes.

#### GLYCINE RECEPTORS ACTIVATION REDUCES $[\text{Cl}^-]_i$ INCREASE BY COUNTERACTING GCM-INDUCED NEURONAL DEPOLARIZATION

Since GCM contains a high concentration of glycine, we hypothesized that GCM-induced  $\text{Cl}^-$  influx, may partially occur through  $\text{Cl}^-$  permeable glycine receptors (GlyR). To test this possibility, we applied GCM to hippocampal neurons together with the GlyR antagonist strychnine ( $1\text{ }\mu\text{M}$ ). Contrary to what expected, we observed an enhancement of GCM-induced  $[\text{Cl}^-]_i$  rise from a control value of  $\Delta R = 0.30 \pm 0.07$  to  $\Delta R = 0.53 \pm 0.12$  ( $n = 9$ ;  $p = 0.05$ , **Figure 4A**). Furthermore, GCM-induced responses in strychnine were completely abolished in the presence of APV and NBQX ( $\Delta R = 0.03 \pm 0.01$ ,  $n = 5$ ). Similar experiments, performed in the presence of gabazine ( $10\text{ }\mu\text{M}$ ), excluded a contribution by GABARs, as no significant changes in  $\Delta R$  were observed ( $0.37 \pm 0.05$  in control vs.  $0.35 \pm 0.08$  in gabazine,  $n = 7$ ; not shown). These data indicate that when GlyRs are blocked, GCM evokes a stronger increase of neuronal  $[\text{Cl}^-]_i$ , entirely dependent on the activation of glutamate receptors.


To investigate whether GCM-induced increase of  $[\text{Cl}^-]_i$  depended on a change in membrane potential, we exposed cell cultures to GCM during current clamp recordings. GCM application (10–15 s) caused the depolarization of the membrane potential of about  $11.4 \pm 3.5\text{ mV}$  ( $n = 27$ , **Figure 4B**), which was absent in the presence of APV and NBQX ( $n = 4$ ; not shown). Conversely, when applied together with strychnine, GCM-induced a significantly higher depolarization ( $33.0 \pm 4.7\text{ mV}$ ,  $p < 0.01$ ;  $n = 20$ ; **Figure 3B**). Thus, the activation of GlyRs, by glycine released from glioma cells, contrasts GCM-induced neuronal depolarization and this effect is abolished when GlyRs are blocked by strychnine. Consistently, in  $\text{Cl}^-$ -Sensor expressing neurons, the reversal potential for chloride was slightly more negative ( $-73 \pm 2\text{ mV}$ ) than the resting membrane potential ( $-66 \pm 4\text{ mV}$ ), as estimated by gramicidin-based perforated patch clamp recordings ( $n = 6$ , not shown), indicating that the activation of GlyRs likely induces neuronal hyperpolarization. Again, GCM-induced depolarization in strychnine was abolished by APV and NBQX ( $2.0 \pm 0.9\text{ mV}$ ,  $n = 7$ , **Figure 4B**), demonstrating that it is entirely dependent on the activation of ionotropic glutamate receptors.

Moreover, when neuronal depolarization was induced by the application of high  $\text{K}^+$  (20 mM, 1 min) an increase in  $[\text{Cl}^-]_i$  rise was induced with a  $\Delta R$  of  $0.29 \pm 0.07$  ( $n = 18$ ,  $p < 0.01$ ; **Figure 4C**), supporting the hypothesis that depolarization is the key event for  $\text{Cl}^-$  entrance into neurons.

#### ANIONIC CHANNELS AND CATIONIC CHLORIDE TRANSPORTERS ARE INVOLVED IN GCM-INDUCED $\text{Cl}^-$ FLUXES

Neuronal depolarization may favor  $\text{Cl}^-$  entry simply augmenting  $\text{Cl}^-$  driving force, or also activating pathways of chloride flux, which are closed at resting potential. Thus we, used a

pharmacological approach trying to shed light on the involvement of  $\text{Cl}^-$  channels and transporters in GCM effects. To investigate the role of transporters in GCM-induced  $\text{Cl}^-$  influx, we exposed hippocampal neurons to the cation chloride transporters blocker furosemide ( $100\text{ }\mu\text{M}$ ). In the presence of this compound,






the application of GCM-evoked a transient increase in  $[\text{Cl}^-]_i$  in average not dissimilar in respect to control ( $R = 0.49 \pm 0.12$  vs.  $0.66 \pm 0.14$ ;  $n = 9$ ;  $p = 0.4$ , paired  $t$ -test; **Figure 5A**). However, in most of the experiments, the amplitude of  $R$  increase was reduced or increased by furosemide, suggesting that both KCC2 and NKCC1 could be involved in GCM-induced  $\text{Cl}^-$  flux. In fact furosemide blocks both transporters, which are expressed in cultured hippocampal neurons. We sought to discriminate between the potentially opposite roles of these two  $\text{Cl}^-$  transporters, by using specific antagonists DIOA and bumetanide.

To block KCC2, we incubated  $\text{Cl}^-$ -Sensor expressing hippocampal cultures in the presence of DIOA ( $20 \mu\text{M}$ ; 2 h), performing experiments in the continuous presence of the drug. In these conditions GCM application still induced an increase of  $R$ , which resulted of  $\Delta R = 0.30 \pm 0.11$  ( $n = 10$ ) compared to slightly higher controls ( $0.55 \pm 0.19$ ,  $n = 8$ ,  $p = 0.25$ , independent  $t$ -test). However, as shown in **Figure 5B**,  $\text{Cl}^-$  did not recover to basal level after GCM washout, indicating that KCC2, although not required for GCM-induced  $\text{Cl}^-$  entry, is necessary for reestablishing  $\text{Cl}^-$  equilibrium. In addition, the basal  $R$  in cells treated with DIOA was higher than control, indicating that prolonged block of KCC2 leads to accumulation of basal  $\text{Cl}^-$

in neurons (see also Pellegrino et al., 2011). Conversely, when we exposed neurons to GCM in the presence of bumetanide, to block NKCC1, we observed an increase in the amplitude of  $R$  transients from a control  $\Delta R$  of  $0.48 \pm 0.13$  to  $0.81 \pm 0.18$  ( $n = 8$ ;  $p = 0.03$ ; **Figure 5C**). This augmented  $\text{Cl}^-$  influx is likely due to the block of NKCC1-mediated rapid  $\text{Cl}^-$  extrusion by reverse transport during intracellular  $\text{Cl}^-$  load (Brumback and Staley, 2008). The block of NKCC1 did not alter the slow recovery after GCM application, indicating that, in our experimental conditions, this transporter is not necessary as KCC2 for the recovery of normal  $\text{Cl}^-$  equilibrium. In addition, the application of bumetanide did not cause significant changes in the basal fluorescence ratio. All together, these data indicate that cation chloride transporters do not play a key role in GCM-induced  $\text{Cl}^-$  entry but their activity may modulate the amount of  $\text{Cl}^-$  entry likely affecting the speed of recovery or changing the driving force of  $\text{Cl}^-$  flux.

It is known that glutamate may lead to neuronal  $\text{Cl}^-$  influx through the activation of different classes of anionic channels (Backus et al., 1998; Van Damme et al., 2003; Slemmer et al., 2004; Inoue and Okada, 2007). Thus, to find out whether anionic channels contributed to GCM-induced neuronal  $\text{Cl}^-$  influx, we tested



**FIGURE 5 |  $\text{Cl}^-$  channels and transporters are involved in**

**GCM-induced  $\text{Cl}^-$  transients. (A)** Effect of furosemide on the peak amplitude of GCM-induced transients ( $n = 9$ ;  $p = 0.4$ , paired  $t$ -test). **(B)** Time course of  $R$  increase induced by GCM in neurons treated with DIOA ( $n = 9$ ,  $\square$ ) and untreated control neurons ( $n = 7$ ,  $\circ$ ). **(C)** Left, example trace representing the effect of acute application of bumetanide on GCM-induced  $\text{Cl}^-$  transient; right, histogram representing the average effect of bumetanide on GCM-induced ratio increase. **(D)** Example trace

of the effect of NPPB ( $40 \mu\text{M}$ ) on GCM-induced  $R$  increase; inset, trace with extended time scale representing GCM application in NPPB.

**(E)** Effect of NPPB  $100 \mu\text{M}$ ; trace and inset as in **(D)**. **(F)** Histogram representing the effect of anionic channels blockers on GCM-induced  $R$  increase in hippocampal neurons. NPPB  $40 \mu\text{M}$  ( $n = 4$ ) and  $100 \mu\text{M}$  ( $n = 5$ ); NFA ( $100 \mu\text{M}$ ;  $n = 3$ ); FFA ( $200 \mu\text{M}$ ;  $n = 5$ ). All data are expressed as the percentage of the control GCM response in the same cells. \* $p < 0.05$ ; \*\* $p < 0.01$ .

the effect of the broad spectrum anion channels blocker 5-Nitro-2-(3-phenylpropylamino) benzoic acid (NPPB, 40–100  $\mu\text{M}$ ).

NPPB, which is known to block a  $\text{Cl}^-$ -driven outwardly rectifying current in neurons (Van Damme et al., 2003), dose dependently reduced GCM-induced increase in  $[\text{Cl}^-]_i$ . At 40  $\mu\text{M}$  it caused a reduction of  $\Delta R$  to  $66 \pm 10\%$  of control ( $n = 4$ ,  $p < 0.05$ ; **Figures 5D,F**) and at 100  $\mu\text{M}$  to  $31 \pm 6\%$  of control ( $n = 9$ ;  $p < 0.01$ ; **Figures 5E,F**), indicating that NPPB-sensitive anion channels largely contribute to GCM-induced  $\text{Cl}^-$  entry in hippocampal neurons. In addition, in the presence of 100  $\mu\text{M}$  NPPB cells were unable to recover  $[\text{Cl}^-]_i$  to basal level after GCM application or neuronal  $\text{Cl}^-$  load. This suggests that anionic channels are involved in GCM-induced  $\text{Cl}^-$  entry, as well as, its recovery. NPPB also caused, by itself, an increase in the basal fluorescence ratio (see **Figures 5D,E**), likely due to  $\text{Cl}^-$  influx, as it was absent when the drug was applied in low external  $\text{Cl}^-$  (8 mM;  $n = 3$ , not shown).

It has been reported a slight reduction effect of NPPB on AMPA-mediated currents (Van Damme et al., 2003). To ascertain the specificity of its effect on  $\text{Cl}^-$  flux, we checked whether this drug could affect the glutamate-induced depolarization in current clamp recordings. Glutamate similarly depolarized neurons in control and in the presence of NPPB, respectively, to  $-10.0 \pm 2.2$  and to  $-13.6 \pm 6.3$  mV ( $n = 5$ ,  $P = 0.5$ ). However, NPPB itself caused a significant depolarization of the neuronal membrane potential of  $10.2 \pm 2.4$  mV ( $n = 8$ , not shown) which could be the cause of NPPB-induced  $\text{Cl}^-$  influx and of the delayed recovery of basal  $\text{Cl}^-$  level.

To further analyze the involvement of anionic channels, we tested the effect of two other anionic channels blockers niflumic acid (NFA) and flufenamic acid (FFA) on GCM response (**Figure 5F**). NFA (100  $\mu\text{M}$ ) reduced the GCM-induced  $\text{Cl}^-$  transient to  $77 \pm 6\%$  of control ( $n = 3$ ;  $p = 0.06$ ), while FFA (200  $\mu\text{M}$ ) was ineffective ( $107 \pm 14$ ;  $n = 5$ ;  $p = 0.78$ ). NFA, similarly to NPPB, also delayed recovery of  $[\text{Cl}^-]_i$  after GCM-induced transients (not shown).

Altogether, these data indicate that the application of GCM mainly determines neuronal  $[\text{Cl}^-]_i$  influx, increasing due to depolarization the driving force for  $\text{Cl}^-$  entry, presumably through anionic channels. Anionic channels and cation chloride transporters are also involved in the recovery of  $[\text{Cl}^-]_i$  equilibrium after GCM-induced  $\text{Cl}^-$  load.

## DISCUSSION

In the present study, we characterized the effect of neuroactive aminoacids released by glioma cells on neuronal  $[\text{Cl}^-]_i$ , taking advantage of a genetically encoded CFP/YFP-based ratiometric  $\text{Cl}^-$ -Sensor (Markova et al., 2008; Bregestovski et al., 2009; Waseem et al., 2010), which was transiently expressed in cultured hippocampal neurons. We observed that acute application of GCM causes a rapid  $[\text{Cl}^-]_i$  increase in neurons via glutamatergic mechanism, mimicked by the application of glutamate and largely dependent on NPPB-sensitive anionic channels. Moreover, HPLC and electrophysiological analysis showed that GCM contains a functionally relevant amount of glutamate and other neuroactive aminoacids, including aspartate, glycine, and serine. Glycine, released by glioma cells, counteracts the

mechanisms triggered by glutamate, likely contrasting neuronal depolarization.

To date, there is little knowledge available on the mechanisms regulating homeostasis of  $\text{Cl}^-$  in either physiological or pathological conditions, especially for the technical difficulties in the measurement of  $\text{Cl}^-$  fluxes. In this paper, we used  $\text{Cl}^-$ -Sensor, a probe with a high sensitivity to  $\text{Cl}^-$  ( $\text{EC}_{50} \sim 30$  mM), that has been demonstrated to be an effective tool for the quantitative estimation of  $[\text{Cl}^-]$  in various cellular compartments (Markova et al., 2008; Bregestovski et al., 2009; Waseem et al., 2010). In our study, several lines of evidence support the notion that GCM alters neuronal  $\text{Cl}^-$  equilibrium, causing glutamatergic-mediated  $\text{Cl}^-$  influx in hippocampal neurons: (i) GCM-induced fluorescence increase is abolished in the presence of antagonists of ionotropic glutamatergic receptors; (ii) exogenously applied glutamate evokes similar  $[\text{Cl}^-]_i$  rise; and (iii) GCM contains glutamatergic agonists at functionally relevant concentration. The ability of glutamate to cause elevation of intracellular  $[\text{Cl}^-]$  has been already shown in physiological (Van Damme et al., 2003; Slemmer et al., 2004; Kitamura et al., 2008) and pathological contexts (Inoue and Okada, 2007; Wang and Qin, 2012). Particularly relevant to our study is that excitotoxic stimulation of glutamate receptors may also lead to pH changes (Wang et al., 1994), since  $\text{Cl}^-$ -Sensor properties might be influenced by acidification (Markova et al., 2008). However, in our experimental conditions, fluorescence ratio increases reasonably represent  $[\text{Cl}^-]_i$  accumulation, because the response evoked by application of glutamate disappeared when we strongly decreased  $\text{Cl}^-$  driving force, lowering external  $[\text{Cl}^-]$  to 8 mM. Consistently, Waseem et al. (2010) and Pellegrino et al. (2011) illustrated the transient glutamate-induced increase of  $[\text{Cl}^-]_i$  in spinal neurons and hippocampal neurons transfected with  $\text{Cl}^-$ -Sensor.

HPLC analysis on GCM confirmed that MZC cells, exposed to control external medium, release a relevant concentration of glutamate, together with other neuroactive aminoacids, giving results very consistent with those found in microdialysis experiments from glioma and peritumoral space in implanted rats (Behrens et al., 2000). In chromatographic analysis we focused on aminoacids commonly involved in signaling mechanisms, although some of them still need to be ascertained (see Behrens et al., 2000). Apparently, the mechanism of glutamate release from glioma cells is different from the most common ways of glutamate transport or release, being insensitive to  $X_c$  system or EAATs inhibition. This is not surprising since  $X_c$  system contribution should be limited in the absence of extracellular cystine and EAATs activity is controversial in glioma cell lines (Ye et al., 1999; Buckingham et al., 2011). Our data with TBOA suggest that EAAT transport system is functional in MZC cells, likely participating in glutamate uptake. Although glutamate release from MZC cells is not evoked by osmotic stress, as described in glial cells, we cannot exclude a contribution of anionic channels, possibly activated by other means (Takano et al., 2005). However, the mechanism of glutamate release by MZC cells needs further investigation. Indeed, other typical ways of astrocytic glutamate release, as P2X7 ATP-gated channels, gap junction hemichannels, or vesicular release (Malarkey and Parpura, 2008), could be functional in glioma cell cultures (see Samadani et al., 2007). Notably, the observed

glutamate release is a specific feature of glioma cells, being glutamate concentration detected in astrocyte conditioned medium, extremely low in comparison to GCM. It should be considered, alternatively, that the observed difference in glutamate concentration might depend on a more functional astrocytic glutamate uptake rather than on different release properties (Takano et al., 2001).

Indeed, the HPLC results showed that GCM contained relevant concentrations of other aminoacids including glycine, serine, and aspartate and might in various ways interfere with neuronal activity. It is known that aminoacids may be released by glial cells both for osmotic (Ordaz et al., 2004) and signaling functions, several of them being already identified as putative gliotransmitter (Behrens et al., 2000; Cavallero et al., 2009). However, some of these substances might have a specific action during glioma invasion and the circumstances of release from glioma cells deserve further investigation.

Our results show that GlyR activation interferes with the depolarizing effect of GCM. The Cl<sup>-</sup> influx through GlyRs may act as a shunting mechanism that counteracts the glutamate-induced depolarization of the membrane potential. Depolarization is likely the key event in GCM-induced Cl<sup>-</sup> influx, as previously proposed for glutamate effects (Backus et al., 1998; Van Damme et al., 2003; Slemmer et al., 2004) and confirmed here by the ability of K<sup>+</sup> to induce a Cl<sup>-</sup> rise similar to that observed with glutamate or GCM. It could be discussed that in the presence of strychnine, GCM-induced Cl<sup>-</sup> accumulation is abolished by GluR antagonists, while a residual Cl<sup>-</sup> transient is observed when GlyRs are not blocked. These results could indicate a participation of GlyRs to Cl<sup>-</sup> influx, particularly in cells with depolarized resting potential.

Cation chloride transporters have a minor role in GCM induced Cl<sup>-</sup> flux, being mainly involved in restoration of Cl<sup>-</sup> equilibrium after GCM stimulation. The potentiating effect of bumetanide is likely due to the block of NKCC1-mediated rapid Cl<sup>-</sup> extrusion by reverse transport during intracellular Cl<sup>-</sup> load (Brumback and Staley, 2008). Alternatively, it is possible to speculate that bumetanide effect on GCM-induced Cl<sup>-</sup> influx, might reflect an increase in Cl<sup>-</sup> driving force, caused by bumetanide mediated decrease in [Cl<sup>-</sup>]<sub>i</sub> (Chabwine et al., 2009). However, this possibility is not supported by our observations.

On the other hand, we show that the depolarization-induced Cl<sup>-</sup> influx involves NPPB-sensitive anionic channels and a significant increase of the neuronal driving force for Cl<sup>-</sup>. The main concern in the interpretation of our results arises from the lack of specific pharmacological tools for anionic channels. In fact, many of the compounds used to block anionic channels are reported to

have unspecific effects and, as shown here, may alter Cl<sup>-</sup> equilibrium and membrane potential. However, our data are strongly consistent with previous reports, showing that anionic channels, activated by depolarization or swelling, are responsible of neuronal Cl<sup>-</sup> entry, following glutamate receptor activation (Van Damme et al., 2003; Slemmer et al., 2004; Inoue and Okada, 2007). We speculate that these may include Ca<sup>2+</sup>-, volume- and voltage-activated Cl<sup>-</sup> channels (Payne et al., 2003; Suzuki et al., 2006; Jentsch, 2008; Deisz et al., 2011). However, other anion permeable channels could be involved, including the high conductance hemichannel Pannexin 1 which is expressed in neurons and is a Cl<sup>-</sup>-permeable and sensitive to NPPB (Ma et al., 2012).

Altogether, our results suggest a mechanism allowing glioma cells to deeply and dynamically interfere with neuronal Cl<sup>-</sup> homeostasis, through the release of excitatory and inhibitory neuroactive substances. In physiological conditions, a similar inhibitory shunt of glutamatergic responses by Cl<sup>-</sup> permeable ligand-gated channels, might occur in case of GABA and glutamate corelease, as described in hippocampal mossy fibers (Treviño et al., 2011), or as a consequence of tonic GABA release (Semyanov et al., 2004). The observed shunt is reminiscent of that caused by unconventional GABA release (Koch and Magnusson, 2009), i.e., during intense synaptic activity (Glykys and Mody, 2007). Interestingly, the resulting excitation/inhibition balance can be altered in epilepsy models, as a results of increased GABAergic inhibition (Treviño et al., 2011). A possible concern in the interpretation of our results is represented by the use of neuronal cultures. In the healthy brain, glutamate is rapidly removed from the extracellular space by astrocytes, preventing extensive or prolonged activation of neuronal glutamate receptors (Danbolt, 2001) and its action may be contrasted by tonic GABA release (Semyanov et al., 2004). Although, it is known that during tumor invasion brain microenvironment is altered (Behrens et al., 2000), the application of GCM might not represent the *natural* concentration profile experienced by neurons. Further *in vivo* studies will be needed to ascertain if the proposed mechanism may support the alterations of Cl<sup>-</sup> homeostasis, causing the imbalance of inhibitory and excitatory neuronal network properties observed in glioma associated epilepsy.

## ACKNOWLEDGMENTS

This paper has been funded by grants from PRIN 2009 (to Davide Ragozzino, Cristina Limatola, and Silvia Di Angelantonio) and Cenci Bolognetti to Cristina Limatola. Emanuele Murana was supported by PhD programme in Neurophysiology, Sapienza University, Rome. Authors wish to thank Dr. Maurizio Barbieri for preliminary chemical determination of CGM cationic content.

## REFERENCES

- Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1999). Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci.* 22, 208–215.
- Backus, K. H., Deitmer, J. W., and Friauf, E. (1998). Glycine-activated currents are changed by coincident membrane depolarization in developing rat auditory brainstem neurones. *J. Physiol.* 507, 783–794.
- Barmashenko, G., Hefft, S., Aertsen, A., Kirschstein, T., and Köhling, R. (2011). Positive shifts of the GABA<sub>A</sub> receptor reversal potential due to altered chloride homeostasis is widespread after status epilepticus. *Epilepsia* 52, 1570–1578.
- Behrens, P. F., Langemann, H., Strohschein, R., Draeger, J., and Hennig, J. (2000). Extracellular glutamate and other metabolites in and around RG2 rat glioma: an intracerebral microdialysis study. *J. Neurooncol.* 47, 11–22.
- Bregestovski, P., Waseem, T., and Mukhtarov, M. (2009). Genetically encoded optical sensors for monitoring of intracellular chloride and chloride-selective channel activity. *Front. Mol. Neurosci.* 2:15. doi: 10.3389/neuro.02.015.2009
- Brumback, A. C., and Staley, K. J. (2008). Thermodynamic regulation of NKCC1-mediated Cl-cotransport underlies plasticity of GABA<sub>A</sub> signaling in neonatal neurons. *J. Neurosci.* 28, 1301–1312.

- Buckingham, S. C., Campbell, S. L., Haas, B. R., Montana, V., Robel, S., Ogunrinu, T., et al. (2011). Glutamate release by primary brain tumors induces epileptic activity. *Nat. Med.* 17, 1269–1274.
- Cavallero, A., Marte, A., and Fedele, E. (2009). L-aspartate as an aminoacid neurotransmitter: mechanisms of the depolarization-induced release from cerebrocortical synaptosomes. *J. Neurochem.* 110, 924–934.
- Chabwine, J. N., Talavera, K., Verbert, L., Eggermont, J., Vanderwinden, J. M., De Smedt, H., et al. (2009). Differential contribution of the Na(+)-K(+)-2Cl(-) cotransporter NKCC1 to chloride handling in rat embryonic dorsal root ganglion neurons and motor neurons. *FASEB J.* 23, 1168–1176.
- Conti, L., Palma, E., Roseti, C., Lauro, C., Cipriani, R., de Groot, M., et al. (2011). Anomalous levels of Cl<sup>-</sup> transporters cause a decrease of GABAergic inhibition in human peritumoral epileptic cortex. *Epilepsia* 52, 1635–1644.
- Danbolt, N. C. (2001). Glutamate uptake. *Prog. Neurobiol.* 65, 1–105.
- Deisz, R. A., Lehmann, T. N., Horn, P., Dehnicke, C., and Nitsch, R. (2011). Components of neuronal chloride transport in rat and human neocortex. *J. Physiol.* 589, 1317–1347.
- De Koninck, Y. (2007). Altered chloride homeostasis in neurological disorders: a new target. *Curr. Opin. Pharmacol.* 7, 93–99.
- Doyon, N., Prescott, S. A., Castonguay, A., Godin, A. G., Kröger, H., and De Koninck, Y. (2011). Efficacy of synaptic inhibition depends on multiple, dynamically interacting mechanisms implicated in chloride homeostasis. *PLoS Comput. Biol.* 7:e1002149. doi: 10.1371/journal.pcbi.1002149
- Fiumelli, H., Cancedda, L., and Poo, M. M. (2005). Modulation of GABAergic transmission by activity via postsynaptic Ca<sup>2+</sup>-dependent regulation of KCC2 function. *Neuron* 48, 773–786.
- Fiumelli, H., and Woodin, M. A. (2007). Role of activity-dependent regulation of neuronal chloride homeostasis in development. *Curr. Opin. Neurobiol.* 17, 81–86.
- Fucile, S., De Saint Jan, D., Prado de Carvalho, L., and Bregestovski, P. (2000). Fast potentiation of glycine receptor channels by intracellular calcium in neurons and transfected cells. *Neuron* 28, 571–583.
- Fucile, S., Milei, R., and Eusebi, F. (2006). Effects of cyclothiazide on GluR1/AMPA receptors. *Proc. Natl. Acad. Sci. U.S.A.* 103, 2943–2947.
- Glykys, J., and Mody, I. (2007). The main source of ambient GABA responsible for tonic inhibition in the mouse hippocampus. *J. Physiol.* 582, 1163–1178.
- Hirschberger, L. L., De La Rosa, J., and Stipanuk, M. H. (1985). Determination of cysteinesulfinate, hypotaurine and taurine in physiological samples by reversed-phase high-performance liquid chromatography. *J. Chromatogr.* 343, 303–313.
- Inoue, H., and Okada, Y. (2007). Roles of volume-sensitive chloride channel in excitotoxic neuronal injury. *J. Neurosci.* 27, 1445–1455.
- Jentsch, T. J. (2008). CLC chloride channels and transporters: from genes to protein structure, pathology and physiology. *Crit. Rev. Biochem. Mol. Biol.* 43, 3–36.
- Kimelberg, H. K., Rutledge, E., Goderie, S., and Charniga, C. (1995). Astrocytic swelling due to hypotonic or high K<sup>+</sup> medium causes inhibition of glutamate and aspartate uptake and increases their release. *J. Cereb. Blood Flow Metab.* 15, 409–416.
- Kitamura, A., Ishibashi, H., Watanabe, M., Takatsuru, Y., Brodwick, M., and Nabekura, J. (2008). Sustained depolarizing shift of the GABA reversal potential by glutamate receptor activation in hippocampal neurons. *Neurosci. Res.* 62, 270–277.
- Koch, U., and Magnusson, A. K. (2009). Unconventional GABA release: mechanisms and function. *Curr. Opin. Neurobiol.* 19, 305–310.
- Lauro, C., Cipriani, R., Catalan, M., Trettel, F., Chece, G., Brusadin, V., et al. (2010). Adenosine A1 receptors and microglial cells mediate CX3CL1-induced protection of hippocampal neurons against Glu-induced death. *Neuropsychopharmacology* 35, 1550–1559.
- Ma, W., Compan, V., Zheng, W., Martin, E., North, R. A., Verkhratsky, A., et al. (2012). Pannexin 1 forms an anion-selective channel. *Pflügers Arch.* 463, 585–592.
- Malarkey, E. B., and Parpura, V. (2008). Mechanisms of glutamate release from astrocytes. *Neurochem. Int.* 52, 142–154.
- Markova, O., Mukhtarov, M., Real, E., Jacob, Y., and Bregestovski, P. (2008). Genetically encoded chloride indicator with improved sensitivity. *J. Neurosci. Methods* 170, 67–76.
- Metzger, F., Repunte-Canonigo, V., Matsushita, S., Akemann, W., Diez-Garcia, J., Ho, C. S., et al. (2002). Transgenic mice expressing a pH and Cl<sup>-</sup> sensing yellow-fluorescent protein under the control of a potassium channel promoter. *Eur. J. Neurosci.* 15, 40–50.
- Ordaz, B., Vaca, L., Franco, R., and Pasantes-Morales, H. (2004). Volume changes and whole cell membrane currents activated during gradual osmolarity decrease in C6 glioma cells: contribution of two types of K<sup>+</sup> channels. *Am. J. Physiol. Cell Physiol.* 286, C1399–C1409.
- O'Shea, R. D. (2002). Roles and regulation of glutamate transporters in the central nervous system. *Clin. Exp. Pharmacol. Physiol.* 29, 1018–1023.
- Palma, E., Amici, M., Sobrero, F., Spinelli, G., Di Angelantonio, S., Ragozzino, D., et al. (2006). Anomalous levels of Cl<sup>-</sup> transporters in the hippocampal subiculum from temporal lobe epilepsy patients make GABA excitatory. *Proc. Natl. Acad. Sci. U.S.A.* 103, 8465–8468.
- Pasantes-Morales, H., Lezama, R. A., Ramos-Mandujano, G., and Tuz, K. L. (2006). Mechanisms of cell volume regulation in hypo-osmolality. *Am. J. Med.* 119, S4–S11.
- Pasantes-Morales, H., and Vázquez-Juárez, E. (2012). Transporters and channels in cytotoxic astrocyte swelling. *Neurochem. Res.* 37, 2379–2387.
- Payne, J. A., Rivera, C., Voipio, J., and Kaila, K. (2003). Cation-chloride co-transporters in neuronal communication, development and trauma. *Trends Neurosci.* 26, 199–206.
- Pellegrino, C., Gubkina, O., Schaefer, M., Becq, H., Ludwig, A., Mukhtarov, M., et al. (2011). Knocking down of the KCC2 in rat hippocampal neurons increases intracellular chloride concentration and compromises neuronal survival. *J. Physiol.* 589, 2475–2496.
- Planells-Cases, R., and Jentsch, T. J. (2009). Chloride channelopathies. *Biochim. Biophys. Acta* 1792, 173–189.
- Rosito, M., DeFlorio, C., Limatola, C., and Trettel, F. (2012). CXCL16 orchestrates adenosine A3 receptor and MCP-1/CCL2 activity to protect neurons from excitotoxic cell death in the CNS. *J. Neurosci.* 32, 3154–3163.
- Samadani, U., Judkins, A. R., Akpalu, A., Aronica, E., and Crino, P. B. (2007). Differential cellular gene expression in ganglioglioma. *Epilepsia* 48, 646–653.
- Sciacaluga, M., Fioretti, B., Catacuzzeno, L., Pagani, F., Bertollini, C., Rosito, M., et al. (2010). CXCL12-induced glioblastoma cell migration require intermediate conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel activity. *Am. J. Physiol. Cell Physiol.* 299, C175–C184.
- Semyanov, A., Walker, M. C., Kullmann, D. M., and Silver, R. A. (2004). Tonically active GABA A receptors: modulating gain and maintaining the tone. *Trends Neurosci.* 27, 262–269.
- Shimamoto, K., Lebrun, B., Yasuda-Kamatani, Y., Sakaitani, M., Shigeri, Y., Yumoto, N., et al. (1998). DL-threo-beta-benzyloxyaspartate, a potent blocker of excitatory aminoacid transporters. *Mol. Pharmacol.* 53, 195–201.
- Slemmer, J. E., Matsushita, S., De Zeeuw, C. I., Weber, J. T., and Knöpfel, T. (2004). Glutamate-induced elevations in intracellular chloride concentration in hippocampal cell cultures derived from EYFP-expressing mice. *Eur. J. Neurosci.* 19, 2915–2922.
- Sontheimer, H. (2003). Malignant gliomas: perverting glutamate and ion homeostasis for selective advantage. *Trends Neurosci.* 26, 543–549.
- Staley, K. J., Soldo, B. L., and Proctor, W. R. (1995). Ionic mechanisms of neuronal excitation by inhibitory GABA receptors. *Science* 269, 977–981.
- Suzuki, M., Morita, T., and Iwamoto, T. (2006). Diversity of Cl<sup>-</sup> channels. *Cell. Mol. Life Sci.* 63, 12–24.
- Takano, T., Kang, J., Jaiswal, J. K., Simon, S. M., Lin, J. H., Yu, Y., et al. (2005). Receptor-mediated glutamate release from volume sensitive channels in astrocytes. *Proc. Natl. Acad. Sci. U.S.A.* 102, 16466–16471.
- Takano, T., Lin, J. H., Arcuino, G., Gao, Q., Yang, J., and Nedergaard, M. (2001). Glutamate release promotes growth of malignant gliomas. *Nat. Med.* 7, 1010–1015.
- Treviño, M., Vivar, C., and Gutierrez, R. (2011). Excitation-inhibition balance in the CA3 network-neuronal specificity and activity-dependent plasticity. *Eur. J. Neurosci.* 33, 1771–1785.
- Van Damme, P., Callewaert, G., Eggermont, J., Robberecht, W., and Van Den Bosch, L. (2003). Chloride influx aggravates Ca<sup>2+</sup>-dependent AMPA receptor-mediated motoneuron death. *J. Neurosci.* 23, 4942–4950.
- Wang, G. J., Randall, R. D., and Thayer, S. A. (1994). Glutamate-induced intracellular acidification of cultured hippocampal neurons



- demonstrates altered energy metabolism resulting from Ca<sup>2+</sup> loads. *J. Neurophysiol.* 72, 2563–2569.
- Wang, Y., and Qin, Z. H. (2012). Molecular and cellular mechanisms of excitotoxic neuronal death. *Apoptosis* 15, 1382–1402.
- Waseem, T., Mukhtarov, M., Buldakova, S., Medina, I., and Bregestovski, P. (2010). Genetically encoded Cl-Sensor as a tool for monitoring of Cl-dependent processes in small neuronal compartments. *J. Neurosci. Methods* 193, 14–23.
- Ye, Z. C., Rothstein, J. D., and Sontheimer, H. (1999). Compromised glutamate transport in human glioma cells: reduction-mislocalization of sodium-dependent glutamate transporters and enhanced. *J. Neurosci.* 19, 10767–10777.
- Ye, Z. C., and Sontheimer, H. (1999). Glioma cells release excitotoxic concentrations of glutamate. *Cancer Res.* 59, 4383–4391.
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Received: 17 September 2012; paper pending published: 16 October 2012; accepted: 06 November 2012; published online: 26 November 2012.
- Citation: Bertollini C, Murana E, Mosca L, D'Erme M, Scala F, Francioso A, Catalano M, Limatola C, Bregestovski P, Di Angelantonio S and Ragozzino D (2012) Transient increase in neuronal chloride concentration by neuroactive aminoacids released from glioma cells. *Front. Mol. Neurosci.* 5:100. doi: 10.3389/fnmol.2012.00100
- Copyright © 2012 Bertollini, Murana, Mosca, D'Erme, Scala, Francioso, Catalano, Limatola, Bregestovski, Di Angelantonio and Ragozzino. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and subject to any copyright notices concerning any third-party graphics etc.